



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	for						Go	Clear
<input checked="" type="checkbox"/> Limits		Preview/Index		History		Clipboard		Details
Display	Abstract	<input type="checkbox"/> Show: 20		<input type="checkbox"/> Sort		Send to		Text

☐ 1: FEBS Lett 1990 Nov 12;274(1-2):23-6

[Related Articles, Links](#)

Entrez
PubMed

Firefly luciferase, synthesized to very high levels in caterpillars infected with a recombinant baculovirus, can also be used as an efficient reporter enzyme in vivo.

Jha PK, Nakhai B, Sridhar P, Talwar GP, Hasnain SE.

PubMed
Services

National Institute of Immunology, Shahid Jeet Singh Marg, New Delhi, India.

Trichoplusia ni and Spodoptera littoralis larvae were infected with a recombinant AcNPV, having the viral polyhedrin gene replaced with the cDNA encoding firefly luciferase. Both S. littoralis and T. ni synthesized very high levels of luciferase representing greater than or equal to 25% and greater than or equal to 15%, respectively of the total Coomassie blue stainable protein. Luciferase was apparently not secreted into the hemolymph but was contained within the body tissue. Expression in S. littoralis larvae suggests that luciferase can be an excellent reporter enzyme to study virus infection, dissemination and expression in different tissues, host range determination, insect physiology and also to monitor the release of recombinant virus in the environment when used as a biocide.

Related
Resources

PMID: 2253776 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="checkbox"/> Show: 20		<input type="checkbox"/> Sort		Send to		Text
---------	----------	-----------------------------------	--	-------------------------------	--	---------	--	------

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

May 29, 2003 11:11 AM